

Supplementary Table 2. Frequency of comedication by classes

	Value
Patients on ≥ 1 comedication	169 (100.0)
Number of comedication per patient	6 (3, 9)
Patients on ≥ 1 comedication, excluding ESRD-associated medications	158 (93.5)
Number of comedication per patient, excluding ESRD-associated medications	4 (2, 6)
Patients on ≥ 1 comedication class	169 (100.0)
Number of comedication classes per patient	3.4 \pm 1.5
Patients on ≥ 1 comedication class, excluding ESRD-associated medications	158 (93.5)
Number of comedication classes per patient, excluding ESRD-associated medications	2.5 \pm 1.5
Number of patients taking comedication in each class	
ESRD-associated medications	159 (94.1)
Cardiovascular agents	118 (69.8)
Anti-diabetic drugs	73 (43.2)
Gastrointestinal agents	62 (36.7)
Central nervous system agent	52 (30.8)
Lipid-lowering agents	36 (21.3)
Anti-microbials	3 (1.8)
Immunosuppressants	4 (2.4)
Other agents	67 (39.6)
Liver protectants	18 (10.7)
Analgesics	16 (9.5)
Medications for lung diseases	15 (8.9)
Urate-lowering drugs	12 (7.1)
Medications for hyperplasia of prostate	9 (5.3)
Anti-histamine	7 (4.1)
Topical medications	5 (3.0)
Medications for thyroid diseases	4 (2.4)
Hormone therapy	1 (0.6)
Other drugs	10 (5.9)

Values are presented as mean \pm standard deviation, median (interquartiles), or number (%).
 ESRD, end-stage renal disease.