

Supplementary Table 3. Treatment responses of DLBCL before and after IPTW

Variables	Before IPTW		After IPTW		P-value
	CHB group (n=82)	Non-CHB group (n=846)	CHB group (n=82)	Non-CHB group (n=846)	
Best response, N (%)					0.02
CR	71 (86.6%)	731 (86.4%)	71 (86.6%)	713 (84.3%)	0.03
PR	4 (4.9%)	39 (4.6%)	4 (4.9%)	55 (6.5%)	
SD	0 (0.0%)	5 (0.6%)	0 (0.0%)	11 (1.3%)	
PD	1 (1.2%)	50 (5.9%)	1 (1.2%)	47 (5.5%)	
Not evaluable	6 (7.3%)	21 (2.5%)	6 (7.3%)	20 (2.4%)	
Objective response rate, N (%)					1.00
CR+PR	75 (91.5%)	770 (91.0%)	75 (91.5%)	770 (91.0%)	0.85
Disease control rate, N (%)					
CR+PR+SD	75 (91.5%)	775 (91.6%)	75 (91.5%)	775 (91.6%)	0.85

CHB, chronic hepatitis B; CR, complete remission; DLBCL, diffuse large B-cell lymphoma; IPTW, inverse probability of treatment weighting; PR, partial response; PD, progressive disease; SD, stable disease.