

Supplementary Table 12. Cause of death among patients with hepatitis C

Rank	Overall (n=4,590)		Male (n=2,804)		Female (n=1,786)	
	Cause of death (KCD code)	Frequency (%)	Cause of death (KCD code)	Frequency (%)	Cause of death (KCD code)	Frequency (%)
1	C22	37.9	C22	41.2	C22	32.7
2	A00-B99	7.5	S00-T98	6.0	A00-B99	11.0
3	S00-T98	5.1	A00-B99	5.2	K74	6.4
4	K74	4.1	C34	3.8	S00-T98	3.7
5	J18	3.0	K74	3.2	J18	3.2
6	C34	2.9	J18	2.9	N18	2.0
7	K70	1.9	K70	2.5	E11	1.8
8	N18	1.9	C25	1.8	C25	1.5
9	E11	1.8	N18	1.8	I21	1.5
10	C25	1.7	E11	1.8	I61	1.5

C22, Malignant neoplasm of liver and intrahepatic bile ducts; A00-B99, Certain infectious and parasitic diseases; S00-T98, Injury, poisoning and certain other consequences of external causes; K74, Fibrosis and cirrhosis of liver; J18, Pneumonia, organism unspecified; C34, Malignant neoplasm of bronchus and lung; K70, Alcoholic liver disease; N18, Chronic kidney disease; E11, Non-insulin-dependent diabetes mellitus; C25, Malignant neoplasm of pancreas; I21, Acute myocardial infarction; I61, Intracerebral hemorrhage.