Supplementary Table 2. The time-dependent areas under receiver operating characteristic curves using the SAFE score or the FIB-4 index for predicting risk of HCC

Subgroups	1-year			3-year			5-year		
	SAFE	FIB-4	*P*-value	SAFE	FIB-4	*P*-value	SAFE	FIB-4	*P*-value
All	0.84	0.83	0.122	0.82	0.82	0.555	0.81	0.80	0.107
HBV	0.81	0.81	0.938	0.77	0.77	0.925	0.79	0.79	0.519
HCV	0.78	0.76	0.312	0.81	0.81	0.816	0.77	0.77	0.444
HBV/HCV	0.91	0.84	<0.001	0.83	0.80	0.223	0.86	0.82	0.207
MASLD	0.36	0.16	<0.001	0.82	0.73	0.047	0.85	0.78	0.008
NAFLD	0.86	0.73	<0.001	0.85	0.71	<0.001	0.88	0.80	0.010
ArLD	NA	NA	NA	0.69	0.67	0.715	0.74	0.69	0.226
Other LC	0.86	0.86	0.997	0.76	0.75	0.669	0.64	0.66	0.531
Other CLD	NA	NA	NA	NA	NA	NA	0.93	0.83	<0.001

SAFE, Steatosis-Associated Fibrosis Estimator; FIB-4, fibrosis-4; HCC, hepatocellular carcinoma; HBV, hepatitis B virus; HCV, hepatitis C virus; MASLD, metabolic dysfunction-associated steatotic liver disease; NAFLD, non-alcoholic fatty liver disease; ArLD, alcohol-related liver disease; LC, liver cirrhosis; Other CLD, other chronic liver disease; NA, not available.

 https://doi.org/10.3350/cmh.2024.0822