특발성 비신단락을 동반한 간경변증 환자의 간성 척수병증 2예

울산대학교 의과대학 서울아산병원 소화기내과학교실, 서울대학교 의과대학 내과학교실 간연구소¹,
서울대학교 의과대학 외과학교실²

구자은 · 임영석 · 명선정¹ · 서경석² · 김강모
이한주 · 정영화 · 이영상 · 서동진

Abstract

Hepatic myelopathy as a presenting neurological complication in patients with cirrhosis and spontaneous splenorenal shunt

Ja Eun Koo, M.D., Young-Suk Lim, M.D., Sun Jeong Myung, M.D.¹,
Kyung Suk Suh, M.D.², Kang Mo Kim, M.D., Han Chu Lee, M.D.,
Young-Hwa Chung, M.D., Yung Sang Lee, M.D., Dong Jin Suh, M.D.

*Department of Gastroenterology, Asan Medical Center,
University of Ulsan College of Medicine, Seoul, Korea*

¹*Department of Internal Medicine and Liver Research Institute,
Seoul National University College of Medicine, Seoul, Korea*

²*Department of Surgery, Seoul National University College of Medicine, Seoul, Korea*

Hepatic myelopathy is a rare complication of chronic liver disease that is associated with extensive portosystemic shunts. The main clinical feature of hepatic myelopathy is progressive spastic paraparesis in the absence of sensory or sphincter impairment. Early and accurate diagnosis of hepatic myelopathy is important because patients with early stages of the disease can fully recover following liver transplantation. Motor-evoked potential studies may be suitable for the early diagnosis of hepatic myelopathy, even in patients with preclinical stages of the disease. Here we describe two patients who presented with spastic paraparesis associated with a spontaneous splenorenal shunt and without any previous episode of hepatic encephalopathy. One patient experienced improved neurologic symptoms after liver transplantation, whereas the other patient only received medical treatment, which did not prevent the progression of spastic paraparesis. (**Korean J Hepatol 2008;14:89-96**)

Key words: Hepatic myelopathy; Liver cirrhosis; Splenorenal shunt

◇ Received March 8, 2007; revised July 29, 2007; accepted September 12, 2007

◇ Abbreviations: ALT, alanine aminotransferase; CT, computed tomography; HIV, human immunodeficiency virus; HM, hepatic myelopathy; INR, international normalized ratio; MEP, motor evoked potential; MR, magnetic resonance; NCS, nerve conduction study; TIPS, transjugular porto-systemic shunt; WBC, white blood cell

◇ Corresponding author: Young-Suk Lim, E-mail: limys@amc.seoul.kr; Phone: 02) 3010-5933; Fax: 02) 3010-5933

주소: 서울시 송파구 풍납동 388-1 서울아산병원 소화기내과 (우) 138-736

INTRODUCTION

Hepatic myelopathy (HM) is a rare neurological complication of chronic liver disease that is characterized clinically by progressive spastic paraparesis without sphincter and sensory impairment.^{1,2} In most reported cases, surgical portosystemic shunts were present during several months to years before symptoms of HM appeared, and HM is rarely found without hepatic encephalopathy.³

This condition of the nervous system is not easily identified, so cases of HM are likely to be overlooked in clinical practice. However, early and accurate diagnosis of HM is important because liver transplantation can be used to treat patients with early stages of this disease, whereas transplantation is less effective for later stages of the disease.^{4,5}

Here, we report two cases of cirrhosis with HM, which was the only neurological complication, and developed after spontaneous splenorenal shunt. One of the patients received a right lobe liver transplant from a living and related donor within 6 months of the appearance of symptoms, and the patient's spastic paraparesis improved slowly over the following 3 years. By contrast, the second patient only received medical treatment aimed to reduce ammonium production and absorption, which did not prevent progression of spastic paraparesis.

Case 1

A 39-year-old man with cirrhosis was referred to our clinic with a walking difficulty which had developed insidiously and progressed for about 3 months. The patient had been diagnosed with hepatitis-B-virus-associated cirrhosis 6

months previously. When admitted to the clinic, he was only able to walk a distance of about 5 meters with the aid of a walking stick. However, he did not have any sphincteric or somatosensory impairment. His liver function was markedly reduced and was classified as Child-Turcotte-Pugh class C, but he had not suffered any hepatic encephalopathy or variceal bleeding. There was no evidence of diabetes mellitus, hypertension, or cardiopulmonary disease. He denied any history of alcohol or illicit drug use, or family history of spastic paraparesis.

Neurologic examination showed the patient to be alert and cranial nerve examination was normal. His muscle tone was increased, particularly in the lower extremities, and he had an ataxic gait. Deep tendon reflex was brisk and plantar reflex showed an extensor response. Lower extremity sensation was normal. There was no evidence of sphincteric, cerebellar, or extrapyramidal dysfunction.

The patient's peripheral blood total white blood cell (WBC) count was 2,500 per mm³ and platelet count was 28,000 per mm³. Prothrombin time was 40.2% (international normalized ratio [INR] of 2.09), serum albumin level was 2.1 g/dL, and total bilirubin was 3.8 mg/dL. Blood ammonia level was increased to 78 µmol/L (normal range is 10~47 µmol/L). Blood electrolyte, urea, creatinine, and alanine aminotransferase (ALT) levels were all in the normal range. A test for serum antibodies to human immunodeficiency virus (HIV) was negative. Due to concerns over thrombocytopenia and impaired coagulation functional status, lumbar puncture was not performed.

A dynamic-enhanced computer tomography (CT) scan of the abdomen revealed cirrhosis of the liver with extensive splenorenal shunt (Fig-

Figure 1. Dynamic-enhanced computer tomography scan of the abdomen shows the extensive splenorenal shunt (arrow).

ure 1). Low-grade esophageal varix was observed during endoscopic examination. Magnetic resonance (MR) images of the cervical, thoracic and lumbar regions of the spine did not reveal any abnormal findings such as cord compression or myelitis (Figure 2). Nerve conduction studies (NCS) were normal.

Considering the clinical findings, this patient was diagnosed as HM associated with hepatitis-B-virus-related cirrhosis with extensive spontaneous splenorenal shunt. A partial liver transplant was performed 3 months later using the right hepatic lobe of a living and related donor. After the transplant, the patient's liver function recovered completely within 4 weeks. There was also slow but progressive improvement in the spastic paraparesis. At the final follow-up visit 3 years after the transplant, the patient was able to walk more than 100 meters unassisted, but still had some difficulty when climbing stairs.

Case 2

A 64-year-old male with cirrhosis due to hepatitis C virus infection was admitted to our

Figure 2. T2-weighted magnetic resonance image of cervico-thoraco-lumbar spine does not show any pathologic lesions.

clinic with a 6-month history of walking difficulty. Weakness of the bilateral lower extremities had developed insidiously and had progressed over several months. When admitted, the patient could only walk about 20 meters unassisted. The patient did not have any difficulty in defecation or voiding, and did not complain of any sensory symptom. In addition, the patient had not previously suffered an episode of hepatic encephalopathy. There was no evidence of diabetes mellitus, hypertension, or cardiopulmonary disease. He denied any history of alcohol, illicit drug use, or family history of spastic paraparesis.

The patient had first been assessed in our clinic 16 months previously for the treatment of a small hepatocellular carcinoma. The tumor was located in the right lobe and was 2.5 cm in diameter. The patient underwent radiofrequency ablation, and there was no evidence of recurrence during the following 2-year period.

Neurologic examination revealed normal mental status and normal cranial nerve function.

Upper-extremity strength was normal. The patient's muscle tone was increased, particularly in the lower extremities with brisk deep-tendon reflex. Plantar reflexes showed an extensor response. Lower-extremity sensation was normal. There was no sphincter dysfunction and no evidence of cerebellar dysfunction or any other extrapyramidal sign.

The peripheral blood total WBC count was 2,800 per mm^3 and the platelet count was 83,000 per mm^3 . Prothrombin time was 68.2% (INR 1.25), albumin level was 3.2 g/dL, and total bilirubin was 2.4 mg/dL. Blood ammonia level was increased to 138 $\mu\text{mol/L}$ (normal range is 10~75 $\mu\text{mol/L}$). Blood electrolyte, urea, creatinine, ALT, vitamin B₁₂, and folate levels were all within normal limits. A test for serum antibodies to HIV was negative. Due to concerns over impaired coagulation status, lumbar puncture was not performed.

Dynamic-enhanced CT of the abdomen revealed cirrhosis of the liver, extensive collateralization, and splenorenal shunt, however, there was no evidence of a recurrence of the hepatocellular carcinoma. MR imaging of the whole spine did not reveal any abnormality, thereby excluding the possibility of cord compression by a metastatic tumor.

The motor-evoked potentials (MEPs) were recorded by transcranial magnetic stimulation studies. Cortical stimulation was performed with a coil connected to a magnetic stimulator, and was applied to the region of the scalp overlying the central sulcus. Magnetic stimulation of the cervical (C7-T1) and lumbar (L4-L5) spinal regions was performed with the same coil. The MEPs were recorded with the electrodes positioned on the abductor digiti minimi muscle of the hand, and on the tibialis anterior muscle of

Figure 3. Motor-evoked potentials (MEPs) after cortical and lumbar/cervical stimulation. MEPs were not obtained from the bilateral tibialis anterior muscles after cortical stimulation, whereas they were found to be normal after lumbar stimulation. In comparison, MEPs recorded from the abductor digiti minimi were normal after both cortical and cervical stimulation. Central motor conduction time (CMCT) was 10.4 ms (normal value ≤ 17 ms).

the leg. Cortical stimulation did not produce MEPs in the bilateral tibialis anterior muscles, whereas lumbar stimulation produced normal MEPs (Figure 3). By contrast, normal MEPs were recorded in the abductor digiti minimi after both cortical and cervical stimulation. These findings indicate motor conduction delay in the dorsal spinal cord.

This patient was diagnosed as HM associated with hepatitis C virus-related cirrhosis and spontaneous extensive splenorenal shunt. He was recommended to receive liver transplant, but a suitable donor could not be found. During the 2-year follow-up period, the patient's hepatic function was conserved and there has been no episode of hepatic encephalopathy. The motor power of his upper extremities and his sphincter function were also conserved. However, his spastic paraparesis has progressed, and at the last follow-up visit he was not able to stand or walk unaided.

Discussion

Hepatic encephalopathy, which is usually reversible, is a common and well-characterized neurologic complication of liver cirrhosis.¹ By contrast, cases of HM that are characterized by progressive spastic paraparesis are rare, and little is known about this condition. HM is irreversible and refractory to many treatments commonly used for hepatic encephalopathy, which aim to lower plasma ammonia levels.⁶ Most neurologists and hepatologists seem to be unaware of this disorder, and it is a frequently overlooked complication in patients with chronic liver disease. Thus, HM is likely to be more frequent than the few case reports indicate.

The diagnosis of HM should be established clinically after exclusion of other possible entities. An MRI scan is necessary to exclude compression of the spinal cord or myelitis. With the exception of an increased glutamine level, evaluation of the cerebrospinal fluid is usually normal.⁷ Specific laboratory tests may be performed to exclude vitamin B₁₂ or folate deficiencies and HIV infection; although these disorders differ markedly from HM in terms of symptoms, and onset, and progression of the disease. The use of an electromyography test to exclude the possibility of second motor neuron involvement often provides little additional information.⁸

The symptoms shown by the patients in this study are typical of HM: spastic paraparesis occurred during the course of liver cirrhosis, which is progressive and permanent. Usually, the onset of HM is insidious and the main clinical feature is spastic paraparesis of the lower extremities.⁹ At first, the symptoms may be slightly asymmetric, but will, as the condition progresses, invariably affect both legs and

steadily worsen. Increased tone and hyperreflexia are the most common findings, and the upper extremities are minimally affected, if at all.¹⁰ Plantar responses are usually extensor, but flexor responses have also been described.^{9,10} Sensorial involvement or sphincteric impairment are uncommon; however, these symptoms have been reported in patients with concurrent neuropathy, probably due to excessive alcohol consumption.^{11,12} Interestingly, almost 90% of the reported cases are in men.³ After several months, progression of the disease plateaus, leaving most patients dependent on assistance or confined to a wheelchair.

Unlike those in previous reports, the patients in this study had not experienced any episodes of hepatic encephalopathy either before or after the occurrence of HM. In most cases, HM symptoms develop after several bouts of hepatic encephalopathy.² Only six patients with HM without previous overt hepatic encephalopathy have been described in the literature.³ In addition, neither patient in this study underwent surgical shunt formation or transjugular portosystemic shunt (TIPS) implantation, but both patients had large spontaneous splenorenal shunts. In most cases, HM symptoms begin after the surgical implantation of a portacaval shunt.³ These findings have led to the assumption that the incidence of HM might decrease as shunt surgery becomes a less common procedure due to the increasing use of endoscopic techniques in the management of esophageal or gastric varices. Nevertheless, cases of HM are still being reported because some cases develop after TIPS implantation and some, like those reported in this study, develop with spontaneous shunting and can be referred to as cases of 'spontaneous hepatic myelopathy'. To our know-

ledge, this study is the first report of patients who developed HM with a spontaneous splenorenal shunt without any episode of hepatic encephalopathy. Therefore, HM must be considered in cirrhotic patients presenting spastic paraparesis without previous hepatic encephalopathy although they have not undergone surgical portacaval shunt or TIPS.

The pathology of HM is known to mainly involve symmetrical demyelination, predominantly of the lateral pyramidal tracts, sometimes associated with axonal loss, and which generally occurs no higher than the cervical cord level.^{9,13} The pathogenesis of HM is poorly understood. However, it is generally accepted that portacaval shunts, or less commonly splenorenal shunts, play an important role,¹¹⁻¹⁵ even in the absence of liver dysfunction. Many authors suggest that shunting of blood may allow nitrogenous breakdown products or neurotoxins to bypass the liver and enter the systemic circulation, resulting in spinal cord damage. Ammonia is the molecule most often thought to be involved.¹¹ Other authors have suggested that hemodynamic factors of the spinal cord may be responsible because the lesions observed are located within spinal segments that do not have an extensive collateral circulation.¹⁴ Nutritional deficiency may also be related to myelopathy.¹⁶ This idea is supported by the fact that pathologic analysis of the spinal cords of prisoners of war and patients suffering from malnutrition are similar to those of patients with HM. Nevertheless, it is not clear why only a few patients with portosystemic shunt develop myelopathy.

Clinical detection is difficult in the early stages of HM and in patients with hepatic encephalopathy or severe peripheral nervous disease. The studies of the MEPs elicited by

transcranial magnetic stimulation have indicated the involvement of the spinal cord in these patients.¹⁷⁻¹⁹ Unfortunately, in the first patient (Case 1), we were unable to analyze MEP generation before and after liver transplantation because of a lack of the required facilities at that time. In Case 2, the MEPs were not recorded from the bilateral tibialis anterior muscles after cortical stimulation, whereas they were shown to be normal after lumbar stimulation. By contrast, MEPs that were recorded from the abductor digiti minimi muscle were normal after both cortical and cervical stimulation (Figure 3). These findings indicate that the lesion is localized between the upper thoracic and lumbar spinal cord and are in agreement with the findings of Nardone et al.,¹⁸ who conducted MEP studies in 13 patients with chronic liver disease associated with portosystemic shunt with or without clinical neurologic symptoms. Nardone et al. confirmed the sensitivity of MEP studies in detecting spinal cord dysfunction, and suggested that MEP studies may show corticospinal electrophysiologic abnormalities even at the preclinical stage of HM, and that these abnormalities become more marked as the disease progresses. Interestingly, they found that patients with early stage disease (as shown by minor neurophysiologic abnormalities) experienced clear neurophysiologic improvement after liver transplantation and an associated marked improvement in neurologic symptoms. By contrast, the clinical and neurophysiologic symptoms of patients with more advanced stages of the disease (shown by marked abnormalities in a MEP study) were not substantially improved by transplantation. Actually, neuropathologic confirmation for HM is very difficult and it is rarely performed. It is supposed that MEP studies for HM are able to

make up for the weak points of clinical diagnosis.

HM is resistant to many therapeutic strategies that are effective in reducing ammonia production and absorption. Two recent studies reported the neurologic outcomes of patients with HM who had undergone liver transplantation.^{4,20} A liver transplant could fully reverse the effects of HM in patients with early-stage disease, whereas it had little effect in patients with more advanced stages of the disease.⁵ These findings indicate that early stages of the disease only involve demyelination of the spinal cord, but that axonal loss occurs at the advanced stage. Thus, early and accurate diagnosis of HM and a subsequent transplantation seem to be important. MEP studies may be useful for the early diagnosis of HM and for monitoring progression of the disease.

In summary, here we describe two patients who developed HM in the presence of spontaneous splenorenal shunt without any episode of hepatic encephalopathy. One patient received medical treatment aimed at reducing ammonia production and absorption, which did not prevent progression of spastic paraparesis. The other patient underwent liver transplantation and exhibited slow but appreciable improvement in neurologic symptoms. Our findings indicate that liver transplantation should be considered as the first therapeutic option of HM and be conducted as early as possible before the spinal cord damage becomes irreversible.

References

1. Jones EA, Weissenborn K. Neurology and the liver. *J Neurol Neurosurg Psychiatry* 1997;63:279-293.
2. Demirci M, Tan E, Elilob B, Gedikoglu G, Saribas O. Spastic paraparesis associated with portal-systemic venous shunting due to congenital hepatic fibrosis. *Neurology* 1992;42:983-985.
3. Conn HO, Rossle M, Levy L, Glocker FX. Portosystemic myelopathy: spastic paraparesis after portosystemic shunting. *Scand J Gastroenterol* 2006;41:619-625.
4. Troisi R, Debruyne J, de Hemptinne B. Improvement of hepatic myelopathy after liver transplantation. *N Engl J Med* 1999;340:151.
5. Counsell C, Warlow C. Failure of presumed hepatic myelopathy to improve after liver transplantation. *J Neurol Neurosurg Psychiatry* 1996;60:590.
6. Campellone JV, Lacomis D, Giuliani MJ, Kroboth FJ. Hepatic myelopathy. Case report with review of the literature. *Clin Neurol Neurosurg* 1996;98:242-246.
7. Sobukawa E, Sakimura K, Hoshino S, Hoshino M, Miyoshi K. Hepatic myelopathy: an unusual neurological complication of advanced hepatic disease. *Intern Med* 1994;33:718-722.
8. Pant SS, Rebeiz JJ, Richardson EP Jr. Spastic paraparesis following portacaval shunts. *Neurology* 1968;18:134-141.
9. Liversedge LA, Rawson MD. Myelopathy in hepatic disease and portosystemic venous anastomosis. *Lancet* 1966;1:277-279.
10. Scobie BA, Summerskill WH. Permanent paraplegia with cirrhosis. *Arch Intern Med* 1964;113:805-810.
11. Bain VG, Bailey RJ, Jhamandas JH. Postshunt myelopathy. *J Clin Gastroenterol* 1991;13:562-564.
12. Lebovics E, DeMatteo RE, Schaffner F, Gendelman S. Portal-systemic myelopathy after portacaval shunt surgery. *Arch Intern Med* 1985;145:1921-1922.
13. Lefer LG, Vogel FS. Encephalomyelopathy with hepatic cirrhosis following portosystemic venous shunts. *Arch Pathol* 1972;93:91-97.
14. Giangaspero F, Dondi C, Scarani P, Zanetti G, Marchesini G. Degeneration of the corticospinal tract following portosystemic shunt associated with spinal cord infarction. *Virchows Arch A Pathol Anat Histopathol* 1985;406:475-481.
15. Dhanwal DK, Das DK. Spastic paraparesis in alcoholic cirrhosis with portal systemic encephalopathy. *J Assoc Physicians India* 1998;46:400.
16. Lemerrier G, Jacquin-Cotton L, Collomb H. Myelopathy of "obscure" etiology--nutritional syndrome. A clinicoanatomic study. *Neurology* 1971;21:185-188.
17. Utku U, Asil T, Balci K, Uzunca I, Celik Y. Hepatic myelopathy with spastic paraparesis. *Clin Neurol Neurosurg* 2005;107:514-516.
18. Nardone R, Buratti T, Oliviero A, Lochmann A, Tezzon F. Corticospinal involvement in patients with a portosystemic shunt due to liver cirrhosis A MEP study. *J Neurol* 2006;253:81-85.
19. Lewis MB, MacQuillan G, Bamford JM, Howdle PD. Delayed myelopathic presentation of the acquired hepatocerebral degeneration syndrome. *Neurology* 2000;54:1011.

20. Weissenborn K, Tietge UJ, Bokemeyer M, Mohammadi B, Bode U, Manns MP, et al. Liver transplantation improves hepatic myelopathy: evidence by three cases. Gastroenterology 2003;124:346-351.